

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : B09B 3/00, C12N 1/00, 5/10, 9/02, 15/53, 15/63, C12P 1/00, 7/02		A1	(11) International Publication Number: WO 99/08812
			(43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/US98/16979		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 17 August 1998 (17.08.98)			
(30) Priority Data: 60/056,754 20 August 1997 (20.08.97) US			
(71) Applicant: THE UNIVERSITY OF ROCHESTER [US/US]; Office of Technology Transfer, 518 Hylan Building, Rochester, NY 14627 (US).			
(72) Inventors: JONES, Jeffrey, P.; 400 N.W. Maryland Court, Pullman, WA 99163 (US). SHIMOJI, Miyuki; Apartment 4C, 229 Conant Road, Rochester, NY 14623 (US).			
(74) Agents: WEYAND, Karla, M. et al.; Nixon, Hargrave, Devans & Doyle LLP, Clinton Square, P.O. Box 1051, P.O. Box 1051, Rochester, NY 14603 (US).		Published <i>With international search report.</i>	
(54) Title: FUNCTIONAL BACTERIAL/MAMMALIAN CYTOCHROME P450 CHIMERA			
(57) Abstract <p>The present invention is directed to a chimeric DNA molecule which includes a first DNA molecule encoding a portion of a full length bacterial P450 protein and a second DNA molecule fused to the first DNA molecule and encoding a portion of a full length mammalian P450 protein. The chimeric DNA molecule encodes a fusion protein which is active and soluble in aqueous liquid. A further aspect of the present invention is directed to the fusion protein encoded by the chimeric DNA molecule. The fusion protein is useful in bioremediation processes and also can be used to hydroxylate a compound to be oxidized.</p>			

FUNCTIONAL BACTERIAL/MAMMALIAN CYTOCHROME P450 CHIMERA

The subject matter of this application was made with support from the
5 United States Government National Institutes of Health Grant No. GM624(PPG),
ES060062, and ES05407. The Government may have certain rights.

This application claims benefit of U.S. Provisional Patent Application
Serial No. 60/056,754, filed August 20, 1997, which is hereby incorporated by
reference.

10

FIELD OF THE INVENTION

The present invention relates to a functional bacterial/mammalian
cytochrome P450 chimera.

15

BACKGROUND OF THE INVENTION

Cytochrome P450 ("P450") is a term used for a widely distributed
group of unique heme proteins which form carbon monoxide complexes with a major
20 absorption band at wavelengths around 450 nm. These proteins are enzymes which
carry out oxidations involved in biosynthesis and catabolism of specific cell or body
components, and in the metabolism of foreign substances entering organisms.
Oxygenating enzymes such as P450 appear to be fundamental cellular constituents in
most forms of aerobic organisms. The activation of molecular oxygen and
25 incorporation of one of its atoms into organic compounds by these enzymes are
reactions of vital importance not only for biosynthesis, but also for metabolic
activation or inactivation of foreign agents such as drugs, food preservatives and
additives, insecticides, carcinogens and environmental pollutants.

In eukaryotic systems P450, and P450 dependent enzymes are known
30 to act on such xenobiotics and pharmaceuticals as phenobarbitol, antipyrine,
haloperidol and prednisone. Known substrates of environmental importance include
compounds such as DDT, and a variety of polychlorinated biphenyls and
polyaromatic hydrocarbons, as well as other halogenated compounds, including
halobenzenes and chloroform.

- 3 -

P4502C9: Baculovirus-mediated Expression, Purification, Structural Characterization, Substrate Stereoselectivity, and Prochiral Selectivity of the Wild-Type and I359L Mutant Forms," Arch. Biochem. Biophys., 333:447-458 (1996); Waterman, M.S., "Heterologous Expression of Mammalian P450 Enzymes," Advances Enzymol., 68:37-66 (1994)) and peptitergents to improve solubility. (Sueyoshi et al., "Molecular Engineering of Microsomal P4502a-4 to a Stable, Water-Soluble Enzyme," Arch. Biochem. Biophys., 322:265-271 (1995)). In contrast, the crystal structures of a number of cytosolic bacterial P450s have been determined. These include P450_{cam}, P450_{bm3}, P450_{terp}, and P450_{eryF}. (Poulos et al., "The 2.6-Å Crystal Structure of *Pseudomonas putida* Cytochrome P-450," J. Biol. Chem., 260:16122-16130 (1985); Poulos et al., "High-Resolution Crystal Structure P450_{cam}," J. Mol. Biol., 195:685-700 (1987); Ravichandran et al., "Crystal Structure of Hemeprotein Domain of P450BM-3, a Prototype for Microsomal P450's," Science, 261:731-736 (1993); Hasemann et al., "Crystal Structure and Refinement of Cytochrome P450_{terp} at 2.3 Å Resolution," J. Mol. Biol., 1169-1185 (1994); Haseman et al., "Structure and Function of Cytochrome P450: A Comparative Analysis of Three Crystal Structures," Structure, 3:41-62 (1995); Cupp-Vickery et al., "Preliminary Crystallographic Analysis of an Enzyme Involved in Erythromycin Biosynthesis: Cytochrome P450_{eryF}," Proteins, 20:197-201 (1994)). Since no detailed structural information has been obtained for a mammalian P450 enzyme, all attempts to determine the effect of enzyme-substrate interactions have used the crystal structures from the soluble bacterial P450 enzymes. (Cupp-Vickery et al., "Preliminary Crystallographic Analysis of an Enzyme Involved in Erythromycin Biosynthesis: Cytochrome P450_{eryF}," Proteins, 20:197-201 (1994); Paulsen et al., Methods in Enzymology, 272:337-46 (1996)). While homology models can be constructed for the membrane-bound mammalian enzymes based on the bacterial enzymes, the very low sequence identities (<20%) mean that any resulting model is of low resolution. In fact, no information directly shows that mammalian and bacterial enzymes are structurally related.

30 The present invention is directed to overcoming the deficiencies of the prior art by forming a P450 protein which is soluble and active in aqueous liquid.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1A is a model of the chimeric structure of the present invention. The blue region is from P450_{cam} and the red region is from CYP2C9. The chimera
 5 contains 3 substrate recognition sites from P450_{cam} and 3 from CYP2C9. Figure 1B shows the construction of a fused plasmid of P450_{cam} and CYP2C9.

Figure 2A is a CO-reduced differential spectrum of the fusion protein of the present invention. The preparation used corresponds to lane 2 in Figure 2B. Figure 2B shows an SDS-polyacrylamide gel electrophoresis of the chimera of the
 10 present invention expressed in *E. coli*. Lanes 1 and 2 show the fusion protein and lane 3 and 4 show P450_{cam} wild-type. Lane 1, 105,000g supernatant (3 µg protein); lane 2, eluate from a hydroxylapatite column (1.5 µg protein); lane 3, 105,000g supernatant (3 µg protein); lane 4, eluate from hydroxylapatite column (2.2 µg protein); lane 5, molecular marker. The gel was stained with Coomassie Brilliant
 15 Blue R250.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a chimeric DNA molecule which
 20 includes a first DNA molecule encoding a portion of a full length bacterial P450 protein and a second DNA molecule fused to the first DNA molecule and encoding a portion of a full length mammalian P450 protein. The chimeric DNA molecule encodes a fusion protein which is active and soluble aqueous liquid. This chimeric DNA molecule can have the nucleotide sequence corresponding to SEQ. ID. No. 1 as
 25 follows:

```

atgacgactg aaaccataca aagcaacgcc aatcttgccc ctctgccacc ccatgtgcc 60
gagcacctgg tattcgactt cgacatgtac aatccgtcga atctgtctgc cggcgtgcag 120
gaggcctggg cagttctgca agaatcaaac gtaccggatc tgggtgtggac tcgctgcaac 180
ggcggacact ggatcgccac tcgcggccaa ctgatccgtg aggcctatga agattaccgc 240
35 cacttttcca gcgagtgtccc gttcatccct cgtgaagccg gcgaagccta cgacttcatt 300
cccacctcga tggatccgcc cgagcagcgc cagtttcgtg cgctggccaa ccaagtgggt 360
ggcatgccgg tgggtggataa gctgggagaac cggatccagg agctggcctg ctccgtgac 420
40 gagagcctgc gcccgcaagg acagtgaac ttcaccgagg actacgccga acccttcccc 480
  
```

- 7 -

Pro Glu Gln Arg Gln Phe Arg Ala Leu Ala Asn Gln Val Val Gly Met
 100 105 110
 5 Pro Val Val Asp Lys Leu Glu Asn Arg Ile Gln Glu Leu Ala Cys Ser
 115 120 125
 Leu Ile Glu Ser Leu Arg Pro Gln Gly Gln Cys Asn Phe Thr Glu Asp
 130 135 140
 10 Tyr Ala Glu Pro Phe Pro Ile Arg Ile Phe Met Leu Leu Ala Gly Leu
 145 150 155 160
 Pro Glu Glu Asp Ile Pro His Leu Lys Tyr Leu Thr Asp Gln Met Thr
 165 170 175
 15 Arg Pro Asp Gly Ser Met Thr Phe Ala Glu Ala Lys Glu Ala Leu Tyr
 180 185 190
 20 Asp Tyr Leu Ile Pro Ile Ile Glu Gln Arg Arg Gln Lys Pro Gly Asn
 195 200 205
 Asn Pro Gln Asp Phe Ile Asp Cys Phe Leu Met Lys Met Glu Lys Glu
 210 215 220
 25 Lys His Asn Gln Pro Ser Glu Phe Thr Ile Glu Ser Leu Glu Asn Thr
 225 230 235 240
 Ala Val Asp Leu Phe Gly Ala Gly Thr Glu Thr Thr Ser Thr Thr Leu
 245 250 255
 30 Arg Tyr Ala Leu Leu Leu Leu Leu Lys His Pro Glu Val Thr Ala Lys
 260 265 270
 35 Val Gln Glu Glu Ile Glu Arg Val Ile Gly Arg Asn Arg Ser Pro Cys
 275 280 285
 Met Gln Asp Arg Ser His Met Pro Tyr Thr Asp Ala Val Val His Glu
 290 295 300
 40 Val Gln Arg Tyr Ile Asp Leu Leu Pro Thr Ser Leu Pro His Ala Val
 305 310 315 320
 Thr Cys Asp Ile Lys Phe Arg Asn Tyr Leu Ile Pro Lys Gly Thr Thr
 325 330 335
 45 Ile Leu Ile Ser Leu Thr Ser Val Leu His Asp Asn Lys Glu Phe Pro
 340 345 350
 50 Asn Pro Glu Met Phe Asp Pro His His Phe Leu Asp Glu Gly Gly Asn
 355 360 365
 Phe Lys Lys Ser Lys Tyr Phe Met Pro Phe Ser Ala Gly Lys Arg Ile
 370 375 380
 55 Cys Val Gly Glu Ala Leu Ala Gly Met Glu Leu Phe Leu Phe Leu Thr
 385 390 395 400
 Ser Ile Leu Gln Asn Phe Asn Leu Lys Ser Leu Val Asp Pro Lys Asn
 405 410 415

- 9 -

Mammalian P450s on Basis of P450_{cam} X-ray Structure," Methods in Enzymology, 206:11-30 (1991), which is hereby incorporated by reference.

Suitable mammalian P450 proteins include 1A, 2B, 2C, 2D, and 3A families of cytochrome P450 and CYP2C9. CYP2C9, which is particularly preferred,
 5 has an amino acid sequence of SEQ. ID. No. 3 as follows:

```

Met Asp Ser Leu Val Val Leu Val Leu Cys Leu Ser Cys Leu Leu Leu
  1           5           10           15
10 Leu Ser Leu Trp Arg Gln Ser Ser Gly Arg Gly Lys Leu Pro Pro Gly
    20           25           30
    Pro Thr Pro Leu Pro Val Ile Gly Asn Ile Leu Gln Ile Gly Ile Lys
      35           40           45
15 Asp Ile Ser Lys Ser Leu Thr Asn Leu Ser Lys Val Tyr Gly Pro Val
    50           55           60
    Phe Thr Leu Tyr Phe Gly Leu Lys Pro Ile Val Val Leu His Gly Tyr
    65           70           75           80
    Glu Ala Val Lys Glu Ala Leu Ile Asp Leu Gly Glu Glu Phe Ser Gly
      85           90           95
25 Arg Gly Ile Phe Pro Leu Ala Glu Arg Ala Asn Arg Gly Phe Gly Ile
    100           105           110
    Val Phe Ser Asn Gly Lys Lys Trp Lys Glu Ile Arg Arg Phe Ser Leu
    115           120           125
30 Met Thr Leu Arg Asn Phe Gly Met Gly Lys Arg Ser Ile Glu Asp Arg
    130           135           140
    Val Gln Glu Glu Ala Arg Cys Leu Val Glu Glu Leu Arg Lys Thr Lys
    145           150           155           160
    Ala Ser Pro Cys Asp Pro Thr Phe Ile Leu Gly Cys Ala Pro Cys Asn
    165           170           175
40 Val Ile Cys Ser Ile Ile Phe His Lys Arg Phe Asp Tyr Lys Asp Gln
    180           185           190
    Gln Phe Leu Asn Leu Met Glu Lys Leu Asn Glu Asn Ile Lys Ile Leu
    195           200           205
45 Ser Ser Pro Trp Ile Gln Ile Cys Asn Asn Phe Ser Pro Ile Ile Asp
    210           215           220
    Tyr Phe Pro Gly Thr His Asn Lys Leu Leu Lys Asn Val Ala Phe Met
    225           230           235           240
    Lys Ser Tyr Ile Leu Glu Lys Val Lys Glu His Gln Glu Ser Met Asp
    245           250           255
  
```

- 11 -

gcatggatat gaagcagtga aggaagccct gattgatctt ggagaggagt tttctggaag 300
 aggcattttc ccactggctg aaagagctaa cagaggattt ggaattgttt tcagcaatgg 360
 5 aaagaaatgg aaggagatcc ggcgtttctc cctcatgacg ctgcggaatt ttgggatggg 420
 gaagaggagc attgaggacc gtgttcaaga ggaagcccg cgccttgagg aggagttgag 480
 10 aaaaaccaag gcctcaccct gtgatccac tttcatcctg ggctgtgctc cctgcaatgt 540
 gatctgctcc attattttcc ataaacgttt tgattataaa gatcagcaat ttcttaactt 600
 aatggaaaag ttgaatgaaa acatcaagat tttgagcagc cctggatcc agatctgcaa 660
 15 taatttttct cctatcattg attacttccc gggaaactcac aacaaattac ttaaaaacgt 720
 tgcttttatg aaaagttata ttttgaaaa agtaaaagaa caccaagaat caatggacat 780
 20 gaacaaccct caggacttta ttgattgctt cctgatgaaa atggagaagg aaaagcacia 840
 ccaaccatct gaatttacta ttgaaagctt ggaaaacact gcagttgact tgtttggagc 900
 tgggacagag acgacaagca caaccctgag atatgctctc cttctcctgc tgaagcacc 960
 25 agaggtcaca gctaaagtcc aggaagagat tgaacgtgtg attggcagaa accggagccc 1020
 ctgcatgcaa gacaggagcc acatgcccta cacagatgct gtggtgcacg aggtccagag 1080
 30 atacattgac cttctcccca ccagcctgcc ccatgcagtg acctgtgaca ttaaattcag 1140
 aaactatctc attcccaagg gcacaacat attaatctcc ctgacttctg tgctacatga 1200
 caacaaagaa tttcccaacc cagagatgtt tgaccctcat cactttctgg atgaaggtgg 1260
 35 caattttaag aaaagtaaat acttcatgcc tttctcagca ggaaaacgga tttgtgtggg 1320
 agaagccctg gccggcatgg agctgttttt attcctgacc tccattttac agaacttta 1380
 40 cctgaaatct ctggttgacc caaagaacct tgacaccact ccagttgtca atggatttgc 1440
 ctctgtgccg cccttctacc agctgtgctt cattcctgct tgaagaagag cagatggcct 1500
 ggctgtgctt gtgcagtccc tgcagctctc tttcctctgg ggcattatcc atctttcact 1560
 45 atctgtaatg ctttttctca cctgtcatct cacattttcc cttccctgaa gatctagtga 1620
 acattcgacc tccattacgg agagtttctt atgtttcact gtgcaaatat atctgctatt 1680
 50 ctccatactc tgtaacagtt gcattgactg tcacataatg ctcatactta tctaagtgtg 1740
 agttattaat atgttattat taaatagaga aatatgattt gtgtattata attcaaaggc 1800
 atttcttttc tgcagtgtct aaataaaaag cattattatt tgctg 1845

55

Suitable bacterial P450 proteins include P450_{cam}, P450_{bm3}, P450_{terp},
 and P450_{eryF}. These proteins are described in Poulos et al., "The 2.6-Å Crystal
 Structure of *Pseudomonas putida* Cytochrome P-450," J. Biol. Chem., 260:16122-
 16130 (1985); Poulos et al., "High-Resolution Crystal Structure P450_{cam}," J. Mol.
 60 Biol., 195:685-700 (1987); Ravichandran et al., "Crystal Structure of Hemeprotein

- 13 -

Asp Ala Ile Ser Ile Val Ala Asn Gly Gln Val Asn Gly Arg Pro Ile
 210 215 220
 5 Thr Ser Asp Glu Ala Lys Arg Met Cys Gly Leu Leu Leu Val Gly Gly
 225 230 235 240
 Leu Asp Thr Val Val Asn Phe Leu Ser Phe Ser Met Glu Phe Leu Ala
 245 250 255
 10 Lys Ser Pro Glu His Arg Gln Glu Leu Ile Glu Arg Pro Glu Arg Ile
 260 265 270
 Pro Ala Ala Cys Glu Glu Leu Leu Arg Arg Phe Ser Leu Val Ala Asp
 275 280 285
 15 Gly Arg Ile Leu Thr Ser Asp Tyr Glu Phe His Gly Val Gln Leu Lys
 290 295 300
 20 Lys Gly Asp Gln Ile Leu Leu Pro Gln Met Leu Ser Gly Leu Asp Glu
 305 310 315 320
 Arg Glu Asn Ala Cys Pro Met His Val Asp Phe Ser Arg Gln Lys Val
 325 330 335
 25 Ser His Thr Thr Phe Gly His Gly Ser His Leu Cys Leu Gly Gln His
 340 345 350
 Leu Ala Arg Arg Glu Ile Ile Val Thr Leu Lys Glu Trp Leu Thr Arg
 355 360 365
 30 Ile Pro Asp Phe Ser Ile Ala Pro Gly Ala Gln Ile Gln His Lys Ser
 370 375 380
 35 Gly Ile Val Ser Gly Val Gln Ala Leu Pro Leu Val Trp Asp Pro Ala
 385 390 395 400
 Thr Thr Lys Ala Val
 405

The DNA molecule encoding P450_{cam} has the nucleotide sequence of
 SEQ. ID. No. 6 as follows:

45 ctgcaggatc gttatccgct ggccgatctg atcaccacgc gtttttccat cgacgaggcc 60
 agcaaggcac ttgaactggt caaggcagga gcaactgatca aaccctgat cgactccact 120
 ctttagccaa cccgcgttcc aggagaacaa caacaatgac gactgaaacc atacaaagca 180
 50 acgccaatct tgccctcttg ccaccccatg tgccagagca cctggtattc gacttcgaca 240
 tgtacaatcc gtcgaatctg tctgcccggc tgcaggaggc ctgggcagtt ctgcaagaat 300
 caaacgtacc ggatctggtg tggactcgct gcaacggcgg aacttgatc gccactcgcg 360
 55 gccaaactgat ccgtgaggcc tatgaagatt accgccactt ttccagcgag tgcccggtta 420
 tccctcgatga agccggcgaa gcctacgact tcattccac ctcgatggat ccgcccagac 480

Natl. Acad. Sci. USA, 268:19681-19689 (1993); Kempf "Truncated Human P450 2D6: Expression in *Escherichia coli*, Ni²⁺-chelate Affinity Purification, and Characterization of Solubility and Aggregation," Arch. Biochem. Biophys., 321:277-288 (1995), which are hereby incorporated by reference).

5 Mutations or variants of the above fusion protein are encompassed by the present invention.

Variants may be modified by, for example, the deletion or addition of amino acids that have minimal influence on the properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated
10 to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification, or identification of the polypeptide.

The DNA molecule encoding the cytochrome P450 polypeptide can be
15 incorporated in cells using conventional recombinant DNA technology. Generally, this involves inserting the DNA molecule into an expression system to which the DNA molecule is heterologous (i.e. not normally present). The heterologous DNA molecule is inserted into the expression system or vector in proper sense orientation and correct reading frame. The vector contains the necessary elements for the
20 transcription and translation of the inserted protein-coding sequences.

U.S. Patent No. 4,237,224 to Cohen and Boyer, which is hereby incorporated by reference, describes the production of expression systems in the form of recombinant plasmids using restriction enzyme cleavage and ligation with DNA ligase. These recombinant plasmids are then introduced by means of transformation
25 and replicated in unicellular cultures including procaryotic organisms and eucaryotic cells grown in tissue culture.

Recombinant genes may also be introduced into viruses, such as vaccinia virus. Recombinant viruses can be generated by transfection of plasmids into cells infected with virus.

30 Suitable vectors include, but are not limited to, the following viral vectors such as lambda vector system gt11, gt WES.tB, Charon 4, and plasmid vectors such as pBR322, pBR325, pACYC177, pACYC184, pUC8, pUC9, pUC18, pUC19,

Similarly, translation of mRNA in procaryotes depends upon the presence of the proper procaryotic signals which differ from those of eucaryotes. Efficient translation of mRNA in procaryotes requires a ribosome binding site called the Shine-Dalgarno ("SD") sequence on the mRNA. This sequence is a short
5 nucleotide sequence of mRNA that is located before the start codon, usually AUG, which encodes the amino-terminal methionine of the protein. The SD sequences are complementary to the 3'-end of the 16S rRNA (ribosomal RNA) and probably promote binding of mRNA to ribosomes by duplexing with the rRNA to allow correct positioning of the ribosome. For a review on maximizing gene expression see
10 Roberts and Lauer, Methods in Enzymology, 68:473 (1979), which is hereby incorporated by reference.

Promoters vary in their "strength" (i.e. their ability to promote transcription). For the purposes of expressing a cloned gene, it is desirable to use strong promoters in order to obtain a high level of transcription and, hence,
15 expression of the gene. Depending upon the host cell system utilized, any one of a number of suitable promoters may be used. For instance, when cloning in *E. coli*, its bacteriophages, or plasmids, promoters such as the T7 phage promoter, *lac* promoter, *trp* promoter, *recA* promoter, ribosomal RNA promoter, the P_R and P_L promoters of coliphage lambda and others, including but not limited, to *lacUV5*, *ompF*, *bla*, *lpp*,
20 and the like, may be used to direct high levels of transcription of adjacent DNA segments. Additionally, a hybrid *trp-lacUV5 (tac)* promoter or other *E. coli* promoters produced by recombinant DNA or other synthetic DNA techniques may be used to provide for transcription of the inserted gene.

Bacterial host cell strains and expression vectors may be chosen which
25 inhibit the action of the promoter unless specifically induced. In certain operons, the addition of specific inducers is necessary for efficient transcription of the inserted DNA. For example, the *lac* operon is induced by the addition of lactose or IPTG (isopropylthio-beta-D-galactoside). A variety of other operons, such as *trp*, *pro*, etc., are under different controls.

30 Specific initiation signals are also required for efficient gene transcription and translation in procaryotic cells. These transcription and translation initiation signals may vary in "strength" as measured by the quantity of gene specific

- 19 -

mismatch will not hybridize (conditions of 85% stringency), and more preferably still, stringent conditions are those under which DNA sequences with more than 10% mismatch will not hybridize (conditions of 90% stringency). In a most preferred embodiment, stringent conditions are those under which DNA sequences with more than 6% mismatch will not hybridize (conditions of 94% stringency).

In yet another aspect of the present invention, the fusion protein can be applied to an environmental pollutant, such as an insecticide or other halogenated hydrocarbon spills, as part of a method of bioremediation. In fact, P450 enzymes can oxidize almost any compound that has a carbon-hydrogen bond and, thus, are useful for almost any environmental contaminant. Generally, microorganisms are extremely useful as agents for clean-up of environmental problems. Development of suitable microorganisms involves either selecting microorganisms with a bioremediation trait or by introducing a gene into microbes to engender them with that ability. By introducing the chimeric DNA molecule into an appropriate vector, it is possible to achieve bioremediation of environmental pollutants. Suitable vectors are non-pathogenic bacteria.

Another aspect of the present invention is using the fusion protein in a process of hydroxylating a compound to be oxidized. Typical compounds to be oxidized include hydrocarbons or any compound having a carbon-hydrogen bond. As discussed above, this involves contacting the compound to be oxidized with the fusion protein under conditions effective to hydroxylate the compound to be oxidized. The fusion protein can be provided by introducing the chimeric DNA molecule into an appropriate vector to express the fusion protein. Suitable vectors include pcW or pkk233-2.

Typicaly, hydroxylation occurs at from about 30 to about 50°C, with 37°C being preferred, with a potassium phosphate buffer and KCl (pH 7.4). The reaction can be monitored by the addition of dichloromethane and assaying by gas chromatography/mass spectrometry.

30

EXAMPLES

The following examples illustrate, but are not intended to limit, the present invention.

- 21 -

primer 7 CATCACCATCACCATCACTGAAGAAGAGCAGATGGCCTGGC
(SEQ. ID. No. 13)

primer 8 GACAGGAATGAAGCACAGCTGGTA (SEQ. ID. No. 14)

5 **Example 2 - Expression of the Fusion Protein**

A single ampicillin-resistant colony of DH5 α cells transformed with plasmid DNA was grown overnight at 37°C in Luria-Bertani medium containing 100 μ g ampicillin ml⁻¹. A 0.5-ml aliquot was used to inoculate 50 ml of Terrific
10 broth ("TB") and cultured for 10 h. This aliquot of 25 ml was used to inoculate 500 ml of TB media. Incubation at 37°C was continued for 19 h. The TB media was supplemented with ampicillin (100 μ g ml⁻¹), 0.2% glucose, 100 μ M δ -aminolevulinic acid, vitamins (100⁻¹w/w, Basal Medium Eagle Vitamin Solution, Gibco BRL, Grand Island, NY), and trace elements (2 mM MgSO₄·7H₂O, 0.1 mM CaCl₂, 1.0 μ M FeSO₄,
15 metal solution1, 50 μ M H₃BO₄, 0.2 μ M CoCl₂·6H₂O, 1 mM CuSO₄·5H₂O, 1 mM MnCl₂·4H₂O, 1 nM Na₂MoO₄ and 2 mM ZnCl₂). The cells were harvested by centrifugation at 5,000 g and 4°C for 10 min. The pellet was stored at -80°C before use.

20 **Example 3 - Construction of Expression Plasmid for Pd and PdR**

Nde I restriction site was introduced at the site of the initiation codon of the Pd or PdR plasmids by the procedures similar to those described above. After digestion of Pd by *Sma* I and digestion of PdR by *Mlu* I followed by blunt-ending,
25 each plasmid was digested by *Nde* I. Gel purified DNA was cloned into PET-15, an expression vector (Novagene, Madison, WI), after digestion by *Xho* I and blunt-ending. *E. coli* strain BL21(DE3) was transformed with pETPd or pETPdR.

Pd and PdR were expressed as follows. Inoculum cultures (25 ml) of *E. coli* BL21(DE3), transformed with pETPd or pETPdR were grown at 37°C in M9
30 minimum medium supplemented with 100 μ g ampicillin ml⁻¹, 0.5% glucose, vitamins, and trace elements as mentioned above. A 25-ml aliquot was used to inoculate 500 ml of M9 minimum medium and the flask was shaken for 1 h at 37°C, at which

Reductase and Puridaredoxin: Cloning, Sequence, and Heterologous Expression of the Proteins," J. Biol. Chem., 265:6066-6073 (1990), which is hereby incorporated by reference). Expression levels of the wild type P450_{cam} was 600-1000 nmoles/liter under similar conditions. After treatment with lysozyme and sonication of the cell pellet, the cell lysate was centrifuged at 105,000g and the supernatant was applied to a Ni-NTA agarose and hydroxylapatite columns (Imai et al., "Expression and Purification of Functional Human 17 α -hydroxylase/17,20-lyase (P45017) in *Escherichia coli*," Proc. Natl. Acad. Sci. USA, 268:19681-19689 (1993), which is hereby incorporated by reference). The purified chimera showed a CO-reduced difference spectrum at 448 nm (Fig. 2A) (Omura et al., "The Carbon Monoxide-Binding Pigment of Liver Microsomes I Evidence for its Hemeprotein Nature," J. Biol. Chem., 239:2370-2378 (1964), which is hereby incorporated by reference), and showed two major bands on SDS-polyacrylamide gel electrophoresis (Fig. 2B) (Laemmli, U.K., "Cleavage of Structural Protein During the Assembly of the Head of Bacteriophage," Nature, 227:680-685 (1970), which is hereby incorporated by reference). Similar bands are observed from purified wild-type P450_{cam} with a [His]₆ tag coding sequence. The lower molecule weight band is presently unidentified. The resulting purified protein showed an approximate molecular weight of 51 kDa as judged by SDS-polyacrylamide gel electrophoresis, consistent with the molecular weight expected for the chimera (Figure 2B).

The resulting purified protein showed a reduced CO difference spectrum at 450 nm (Figure 2A). These data are consistent with a folded P450 protein having a functional active site. The observation that a functional chimera of P450 2C9 and P450_{cam}, which have only 15% primary sequence homology, can still bind CO provides strong evidence for a conserved three-dimensional structure between P450_{cam} and CYP2 family. The fact that the resulting enzyme is soluble, while mammalian enzymes with the amino terminus removed are not, indicates that other regions near the amino terminus may also be important for membrane interactions. (Lemos-Chiarandine et al., J. Cell Biol., 104:209-219 (1987); Vergeres et al., Biochemistry, 28:3650-3655 (1989); Wachenfeldt et al., Arch. Biochem. Biophys., 339:107-114 (1997), which are hereby incorporated by reference.)

- 25 -

putidaredoxin reductase, and 300 μ M NADH. The reaction was stopped by the addition of 4 ml of dichloromethane and assayed by gas chromatography/mass spectrometry. Experiments to determine if the mammalian P450 reductase can support the same oxidation are underway.

5 Detection of the catalytic activity toward 4-chlorotoluene indicate that the fusion protein can function as an active P450 enzyme (Table 1). As compared with the turnover number from the wild type P450_{cam}, the chimera shows approximately 3 times the activity towards 4-chlorotoluene. This means a potential for making soluble P450 that can perform stereospecific synthesis.

10 This approach could have a number of applications. 1) From other homology models of mammalian P450 enzymes it is apparent that this method may prove to be a general method for constructed soluble P450 enzymes with mammalian active site characteristics. These enzymes should be more adaptable to uses in benign synthesis and bioremediation than the more restrictive bacterial enzymes and easier to
15 work with then the membrane bound mammalian enzymes. 2) Selectively replacing amino acid segments in the amino terminus with the mammalian amino acids may prove to be a valuable method of determining important membrane association sites. 3) Since the enzyme is soluble, it could prove a method for obtaining structural information. In particular it should be amiable to Xray crystallography. 4) Since the
20 enzyme is part mammalian and part bacterial, it can be used to determine the features that confer specific interactions with the different reductases system that are used by the bacterial and mammalian proteins.

 Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is solely for that purpose, and variations
25 can be made therein by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

8. A chimeric DNA molecule according to claim 1, wherein the chimeric DNA molecule has a heme ligand positioned in a relative orientation to an I-helix and a fifth cysteine ligand similar to that of the heme ligand in a full length mammalian P450 protein.
9. A chimeric DNA molecule according to claim 1, wherein the chimeric DNA molecule encodes an amino acid sequence of SEQ. ID. No. 2.
10. A chimeric DNA molecule according to claim 9, wherein the chimeric DNA molecule has a nucleotide sequence of SEQ. ID. No. 1.
11. A DNA expression system transformed with the chimeric DNA molecule of claim 1.
12. A DNA expression system according to claim 11, wherein the chimeric DNA molecule is positioned in the expression system in proper sense orientation and correct reading frame.
13. A DNA expression system according to claim 11, wherein the first and second DNA molecules are fused together at a location where the encoded fusion protein lacks secondary structure.
14. A host cell transformed with the chimeric DNA molecule of claim 1.
15. A host cell according to claim 14, wherein the host cell is selected from the group consisting of plant cells, mammalian cells, insect cells, and bacterial cells.
16. A fusion protein comprising:
a portion of a bacterial P450 protein and
a portion of a mammalian P450 protein fused to the portion of a bacterial P450 protein, wherein the fusion protein is active and soluble in aqueous liquid.

- 29 -

contacting the compound to be oxidized with the fusion protein according to claim 16 under conditions effective to hydroxylate the compound to be oxidized.

5 26. A method according to claim 25, wherein the portion of the mammalian P450 protein and the portion of the bacterial P450 protein are fused where the encoded fusion protein lacks secondary structure.

10 27. A method according to claim 25, wherein the fusion protein is prepared from a full length mammalian P450 protein where a portion of the full length mammalian P450 protein is replaced with a homologous portion of a full length bacterial P450 protein.

15 28. A method according to claim 27, wherein all amino acids prior to a random coil between G- and H-helices in the full length mammalian P450 protein are replaced with a homologous portion of the full length bacterial P450 protein.

20 29. A method according to claim 27, wherein the fusion protein comprises about 50 percent of the full length mammalian P450 protein and about 50 percent of the full length bacterial P450 protein.

 30. A method according to claim 25, wherein the fusion protein is provided by providing a vector comprising a chimeric DNA molecule comprising:
 a first DNA molecule encoding a portion of a full length
25 bacterial P450 protein;


 a second DNA molecule fused to the first DNA molecule and encoding a portion of a full length mammalian P450 protein, wherein the chimeric DNA molecule encodes the fusion protein.

30 31. A method according to claim 30, wherein the first and second DNA molecules are fused together at a location where the encoded fusion protein lacks secondary structure.

 32. A method according to claim 30, wherein the chimeric DNA
35 molecule is prepared from a DNA molecule encoding a full length mammalian P450

- 31 -

40. A method according to claim 35, wherein the fusion protein is provided by providing a vector comprising a chimeric DNA molecule comprising:
a first DNA molecule encoding a portion of a full length bacterial P450 protein;
- 5 a second DNA molecule fused to the first DNA molecule and encoding a portion of a full length mammalian P450 protein, wherein the chimeric DNA molecule encodes the fusion protein.
41. A method according to claim 40, wherein the first and second
10 DNA molecules are fused together at a location where the encoded fusion protein lacks secondary structure.
42. A method according to claim 40, wherein the chimeric DNA molecule is prepared from a DNA molecule encoding a full length mammalian P450
15 protein where a portion of the DNA molecule encoding a full length mammalian P450 protein is replaced with a DNA molecule encoding a homologous portion of a full length bacterial P450 protein.
43. A method according to claim 42, wherein all amino acids prior
20 to a random coil between G- and H-helices in the full length mammalian P450 protein are replaced with a homologous portion of the full length bacterial P450 protein.
44. A method according to claim 42, wherein the chimeric DNA molecule comprises about 50 percent of the DNA molecule encoding the full length
25 mammalian P450 protein and about 50 percent of the DNA molecule encoding the full length bacterial P450 protein.



4/4




FIG. 2B

ctggccggca tggagctgtt tttattcctg acctccattt tacagaactt taacctgaaa 1260
 tctctggttg acccaaagaa ccttgacacc actccagttg tcaatggatt tgcctctgtg 1320
 ccgcccttct accagctgtg cttcattcct gtctga 1356

<210> 2

<211> 446

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: fusion protein

<400> 2

Asn Leu Ala Pro Leu Pro Pro His Val Pro Glu His Leu Val Phe Asp
 1 5 10 15

Phe Asp Met Tyr Asn Pro Ser Asn Leu Ser Ala Gly Val Gln Glu Ala
 20 25 30

Trp Ala Val Leu Gln Glu Ser Asn Val Pro Asp Leu Val Trp Thr Arg
 35 40 45

Cys Asn Gly Gly His Trp Ile Ala Thr Arg Gly Gln Leu Ile Arg Glu
 50 55 60

Ala Tyr Glu Asp Tyr Arg His Phe Ser Ser Glu Cys Pro Phe Ile Pro
 65 70 75 80

Arg Glu Ala Gly Glu Ala Tyr Asp Phe Ile Pro Thr Ser Met Asp Pro
 85 90 95

Pro Glu Gln Arg Gln Phe Arg Ala Leu Ala Asn Gln Val Val Gly Met
 100 105 110

Pro Val Val Asp Lys Leu Glu Asn Arg Ile Gln Glu Leu Ala Cys Ser
 115 120 125

Leu Ile Glu Ser Leu Arg Pro Gln Gly Gln Cys Asn Phe Thr Glu Asp
 130 135 140

Tyr Ala Glu Pro Phe Pro Ile Arg Ile Phe Met Leu Leu Ala Gly Leu
 145 150 155 160

Pro Glu Glu Asp Ile Pro His Leu Lys Tyr Leu Thr Asp Gln Met Thr
 165 170 175

Arg Pro Asp Gly Ser Met Thr Phe Ala Glu Ala Lys Glu Ala Leu Tyr
 180 185 190

<210> 3

<211> 490

<212> PRT

<213> mammalian

<400> 3

Met Asp Ser Leu Val Val Leu Val Leu Cys Leu Ser Cys Leu Leu Leu
 1 5 10 15

Leu Ser Leu Trp Arg Gln Ser Ser Gly Arg Gly Lys Leu Pro Pro Gly
 20 25 30

Pro Thr Pro Leu Pro Val Ile Gly Asn Ile Leu Gln Ile Gly Ile Lys
 35 40 45

Asp Ile Ser Lys Ser Leu Thr Asn Leu Ser Lys Val Tyr Gly Pro Val
 50 55 60

Phe Thr Leu Tyr Phe Gly Leu Lys Pro Ile Val Val Leu His Gly Tyr
 65 70 75 80

Glu Ala Val Lys Glu Ala Leu Ile Asp Leu Gly Glu Glu Phe Ser Gly
 85 90 95

Arg Gly Ile Phe Pro Leu Ala Glu Arg Ala Asn Arg Gly Phe Gly Ile
 100 105 110

Val Phe Ser Asn Gly Lys Lys Trp Lys Glu Ile Arg Arg Phe Ser Leu
 115 120 125

Met Thr Leu Arg Asn Phe Gly Met Gly Lys Arg Ser Ile Glu Asp Arg
 130 135 140

Val Gln Glu Glu Ala Arg Cys Leu Val Glu Glu Leu Arg Lys Thr Lys
 145 150 155 160

Ala Ser Pro Cys Asp Pro Thr Phe Ile Leu Gly Cys Ala Pro Cys Asn
 165 170 175

Val Ile Cys Ser Ile Ile Phe His Lys Arg Phe Asp Tyr Lys Asp Gln
 180 185 190

Gln Phe Leu Asn Leu Met Glu Lys Leu Asn Glu Asn Ile Lys Ile Leu
 195 200 205

Ser Ser Pro Trp Ile Gln Ile Cys Asn Asn Phe Ser Pro Ile Ile Asp

465

470

475

480

Pro Phe Tyr Gln Leu Cys Phe Ile Pro Val

485

490

<210> 4

<211> 1845

<212> DNA

<213> mammalian

<400> 4

```

gaaggttca atggattctc ttgtggctct tgtgctctgt ctctcatgtt tgcttctcct 60
ttcactctgg agacagagct ctgggagagg aaaactccct cctggcccca ctctctctcc 120
agtgattgga aatatacctac agataggtat taaggacatc agcaaatacct taaccaatct 180
ctcaaaggtc tatggccctg tgttcaactct gtattttggc ctgaaaccca tagtggtgct 240
gcatggatat gaagcagtga aggaagccct gattgatctt ggagaggagt tttctggaag 300
aggcattttc cactggctg aaagagctaa cagaggattt ggaattgttt tcagcaatgg 360
aaagaaatgg aaggagatcc ggcgtttctc cctcatgacg ctgcggaatt ttgggatggg 420
gaagaggagc attgaggacc gtgttcaaga ggaagcccg cgccttgtgg aggagttgag 480
aaaaaccaag gcctcaccct gtgatccac tttcatcctg ggctgtgctc cctgcaatgt 540
gatctgctcc attattttcc ataaacgttt tgattataaa gatcagcaat ttcttaactt 600
aatggaaaag ttgaatgaaa acatcaagat tttagcagc ccctggatcc agatctgcaa 660
taatttttct cctatcattg attacttccc gggaactcac aacaaattac ttaaaaacgt 720
tgcttttatg aaaagttata ttttgaaaaa agtaaaagaa caccaagaat caatggacat 780
gaacaaccct caggacttta ttgattgctt cctgatgaaa atggagaagg aaaagcacaa 840
ccaaccatct gaatttacta ttgaaagctt ggaaaacact gcagttgact tgtttggagc 900
tgggacagag acgacaagca caaccctgag atatgctctc cttctcctgc tgaagcacc 960
agaggtcaca gctaaagtcc aggaagagat tgaacgtgtg attggcagaa accggagccc 1020
ctgcatgcaa gacaggagcc acatgcccta cacagatgct gtggtgcacg aggtccagag 1080
atacattgac cttctcccca ccagcctgcc ccatgcagtg acctgtgaca ttaaattcag 1140
aaactatctc attcccaagg gcacaacatc attaatctcc ctgacttctg tgctacatga 1200
caacaaagaa tttcccaacc cagagatgtt tgaccctcat cactttctgg atgaaggagg 1260
caatttttaag aaaagtaaat acttcatgcc tttctcagca ggaaaacgga tttgtgtggg 1320
agaagccctg gccggcatgg agctgttttt attcctgacc tccattttac agaacttta 1380
cctgaaatct ctggttgacc caaagaacct tgacaccact ccagttgtca atggatttgc 1440
ctctgtgccg ccttctacc agctgtgctt cattcctgtc tgaagaagag cagatggcct 1500
ggctgctgct gtgcagtccc tgcagctctc tttcctctgg ggcattatcc atctttcact 1560
atctgtaatg cttttctca cctgtcatct cacattttcc cttccctgaa gatctagtga 1620
acattcgacc tccattacgg agagtttctt atgtttcact gtgcaaatat atctgctatt 1680
ctccatactc tgtaacagtt gcattgactg tcacataatg ctcatactta tctaattgtt 1740
agttattaat atgttattat taaatagaga aatatgattt gtgtattata attcaaaggc 1800
atttcttttc tgcattgtct aaataaaaag cattattatt tgctg 1845

```

<210> 5

<211> 405

<212> PRT

<213> Pseudomonas putida

245 250 255
 Lys Ser Pro Glu His Arg Gln Glu Leu Ile Glu Arg Pro Glu Arg Ile
 260 265 270
 Pro Ala Ala Cys Glu Glu Leu Leu Arg Arg Phe Ser Leu Val Ala Asp
 275 280 285
 Gly Arg Ile Leu Thr Ser Asp Tyr Glu Phe His Gly Val Gln Leu Lys
 290 295 300
 Lys Gly Asp Gln Ile Leu Leu Pro Gln Met Leu Ser Gly Leu Asp Glu
 305 310 315 320
 Arg Glu Asn Ala Cys Pro Met His Val Asp Phe Ser Arg Gln Lys Val
 325 330 335
 Ser His Thr Thr Phe Gly His Gly Ser His Leu Cys Leu Gly Gln His
 340 345 350
 Leu Ala Arg Arg Glu Ile Ile Val Thr Leu Lys Glu Trp Leu Thr Arg
 355 360 365
 Ile Pro Asp Phe Ser Ile Ala Pro Gly Ala Gln Ile Gln His Lys Ser
 370 375 380
 Gly Ile Val Ser Gly Val Gln Ala Leu Pro Leu Val Trp Asp Pro Ala
 385 390 395 400
 Thr Thr Lys Ala Val
 405

<210> 6

<211> 1578

<212> DNA

<213> *Pseudomonas putida*

<400> 6

ctgcaggatc gttatccgct ggccgatctg atcaccacgc gtttttccat cgacgaggcc 60
 agcaaggcac ttgaactggt caaggcagga gcaactgatca aaccctgat cgactccact 120
 ctttagccaa cccgcgttcc aggagaacaa caacaatgac gactgaaacc atacaaagca 180
 acgccaatct tgcccctctg ccaccccatg tgccagagca cctggtattc gacttcgaca 240
 tgtacaatcc gtcgaatctg tctgcggcg tgccaggagc ctgggcagtt ctgcaagaat 300
 caaacgtacc ggatctggtg tggactcgct gcaacggcgg aacttgatc gccactcgcg 360
 gccaaactgat ccgtgaggcc tatgaagatt accgccactt ttccagcgag tgcccgttca 420
 tccctcgtga agccggcgaa gcctacgact tcattccac ctcgatggat ccgcccgcgc 480
 agcgccagtt tcgtgcgctg gccaaccaag tggttggcat gccggtggtg gataagctgg 540

<400> 9
ccatggacaa ccctcaggac tttattgat 29

<210> 10
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic

<400> 10
ccattgattc ttggtgttct tttact 26

<210> 11
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic

<400> 11
gcatgaacaa ccctcaggac tttattga 28

<210> 12
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic

<400> 12
ccggcttctg cctgcgttgc tcg 23

<210> 13
<211> 41
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:synthetic

<400> 13
catcaccatc accatcactg aagaagagca gatggcctgg c 41

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16979

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : B09B 3/00; C12N 1/00, 5/10, 9/02, 15/53; 15/63; C12P 1/00, 7/02

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.2; 23.7; 435/41, 56, 57, 58, 59, 61, 125, 189, 262.5, 69.1, 320.1, 252.3, 254.11, 325, 410

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,114,852 A (YABUSAKI et al.) 19 May 1992, entire document.	1-44
A	US 5,240,831 A, (H.J. BARNES) 31 August 1993, entire document.	1-44
A	O'KEEFE et al. Occurrence and biological function of cytochrome P450 monooxygenases in the actinomycetes. Molecular Microbiology. 1991. Vol. 5, No. 9, pages 2099-2105, entire document.	25-44
A	OKUDA et al. Recent progress in enzymology and molecular biology of enzymes involved in vitamin D metabolism. Journal of Lipid Research. 1995. Vol. 36, pages 1641-1652, entire document.	25-34

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 SEPTEMBER 1998

Date of mailing of the international search report

23 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GABRIELE E. BUGAISKY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16979

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

536/23.2; 435/41, 189, 262.5, 69.1, 320.1, 252.3, 254.11, 325, 410

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, STN-CAS files Registry, Caplus, Biotechds, Derwent WPI; A-geneseq32, pir56, swissprot35, sptrembl16
search terms: cytochrome p450, fusi?, chimer?, bacter?, prokaryot?, eukaryot?, yeast mammalian, pseudomonas, putida,
cyp2c9